



Cool Season Lawn Calendar—Eastern Nebraska

Sarah Browning, University of Nebraska Extension Educator

John Fech, University of Nebraska Extension Educator

Amanda Folck, University of Nebraska Turfgrass Extension Educator, Agronomy and Horticulture

Roch Gaussoin, University of Nebraska Turfgrass Specialist, Agronomy and Horticulture

A complicating factor in today's garden market is turfgrass fertilizers and preemergence weed control products are often available only as combined products. Newer (less than 20 years old) and older lawns should be managed differently for resource efficiency and environmental stewardship. This publication provides timely information for Kentucky bluegrass, tall fescue and other cool season grasses (perennial ryegrass and fine fescues.)

Newer lawns often struggle due to soil low in nutritional value and high weed pressure. They are best managed by making a fertilizer plus preemergence herbicide application in late April/early May followed by a second application in early June.

Older lawns need less fertilization to perform well and have lower weed pressure. Preemergence weed control plus fertilizer applications are optional based on need and desired outcome.

RESOURCES

- Proper Lawn Aeration—<https://bit.ly/CSUaeration>
- Bluegrass Billbug—<https://bit.ly/UMNbillbug>
- Managing Billbugs in Turfgrass—<https://bit.ly/PUbillbug>
- Broadleaf weed control—<https://go.unl.edu/NEbroadleafweed>
- Brown Patch—<https://byf.unl.edu/brown-patch>
- Crabgrass control—<https://go.unl.edu/NEcrabgrass>
- Establishing Lawns from Seed—<https://go.unl.edu/NETurf-seed>
- Improving Turf in Fall—<https://go.unl.edu/NETurf-fall>
- Irrigation audit—<https://go.unl.edu/NEirrigation-audit>
- Summer Patch—<https://bit.ly/PSUsummer-patch>
- White Grubs—<https://bit.ly/PUwhite-grub>

This publication has been peer reviewed.
Nebraska Extension publications are available
online at <http://extensionpubs.unl.edu/>.

Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska-Lincoln cooperating with the Counties and the United States Department of Agriculture.

Nebraska Extension educational programs abide with the nondiscrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.

© 2023, The Board of Regents of the University of Nebraska on behalf of the Nebraska Extension.
All rights reserved.

Fertilization Dates	Newer Lawn	Older Lawn	Cultural Practices	Pest Control	Notes
April					
April 25-May 1	0.75—1 lbs N/1000 sq.ft.	⁰ Optional - 0.25 lbs N/1000 sq.ft.	Aerate ¹ Summer patch prevention ²	Billbugs ³ Preemergence weed control—1 st application ⁴	Mow at 3–3.5 inches all summer. Avoid removing more than 1/3 leaf height at one time. ⁰ If turf characteristics, such as color and density, are not ideal in early season, fertilization can be made. Otherwise, it can be skipped. ¹ Use hollow tines for maximum compaction reduction. ² When 2-inch soil depth reaches 65F temp., apply fungicides to lawns with history of infection. Repeat application 4 weeks later. ³ If lawn has a history of billbug damage. Look for adult feeding symptoms or activity in full sun near sidewalks. ⁴ When a 4-inch soil depth reaches 55F temp, apply fertilizer plus preemergence herbicide for annual weed control, including crabgrass and foxtail.
April 25-May 15			Overseed ⁵		⁵ Core aerify before or power rake after seeding for good seed-soil contact. With new seedings, use only a preemergence products containing mesotrione or topramezone to prevent seedling death.
June 1–15	0.5 lbs N/1000 sq.ft.	0.5 lbs N/1000 sq.ft.		Preemergence weed control—2 nd application	
June–Sept.	Irrigate to prevent drought stress		Irrigation/sprinkler audit ⁶	Scout for dollar spot and brown patch. ⁷	⁶ Check uniformity of sprinkler output and equipment function. Turf requires 1" water/week for good growth, including both rain and irrigation. Overwatering weakens the turf. ⁷ Fungicides are not usually recommended for home lawns except in extreme cases.
June 25-July 4				White grub control	Apply preventive insecticides if lawn has a history of white grub damage.
Aug. 15–Sept. 15			Overseed ⁵		
Aug. 25–Sept. 5	0.75—1 lbs N/1000 sq.ft.				
Sept 1-Oct 15				Winter annual weed control.	For lawns with a history of winter annual weeds, such as henbit and annual bluegrass, apply a fall preemergence herbicide to kill germinating seeds.
Sept 15				0.5–1 lb.	One fall application with 30–50% quick release/water soluble nitrogen. This will provide even release throughout fall.
Sept. 20-Oct. 20				Aerate ¹	Spot spray as needed. Fall is ideal time to control broadleaf weeds.
Oct. 15–30	0.5 lbs N/1000 sq.ft.	Optional - 0.25 lbs N/1000 sq.ft.	Broadleaf perennial weed control		Second best time is in spring shortly after flowering of dandelions. Only needed on new lawns or areas that are thin or have suffered pest damage.
Oct. 15-Nov. 1					Continue mowing until grass stops growing.