



Creep Feeding Beef Calves

Rick Rasby, Extension Specialist

Kacie McCarthy, Extension Beef Cow-Calf Specialist

Management and creep feeding calves are covered in this NebGuide.

The definition of creep feeding in this NebGuide is providing the calf with a high-quality feed resource, free choice, in addition to milk nursed from the dam and any other forage resource that the calf has access to consume.

Gross income of the cow-calf enterprise is largely dependent on weaning weight of the calves. There are strategies that can increase calf weaning weight if weaning date is not changed: increase milk production of the dam; selection pressure to growth; increase forage consumption of the calf (depends on forage quality); or provide supplemental feed to the calf that is high quality and nutrient dense.

Management practices exist to provide calves with the highest-quality standing forage (creep grazing) but will require labor and fencing infrastructure. Increasing milk production of the dam will increase weaning weight but will also increase the nutrient requirements of the dam, increase daily feed intake and will reduce the number of cows that can graze a fixed forage resource base.

Creep feeding studies consistently show an increase in weaning weight. Creep feeding beef calves is usually reserved for certain market/management situations such as high calf prices in conjunction with low feed prices, dry lot cow-calf operations, fall-born calves, possibly in drought conditions to sustain minimal calf growth, and seedstock

operations. Because data suggest non-creep fed calves catch up weight-wise with their creep-fed mates post-weaning, the greatest return is realized if calves are sold at weaning.

Under severe drought conditions, creep feeding can be used to sustain minimal calf growth. A risk of creep feeding is getting calves too fat, resulting in price discounts or lowered lifetime milk production of heifer calves intended to become replacements for the cow herd.

Creep feeding must be carefully appraised in view of the economics of cost, potential market, and the influence on sale price of calves.

During drought conditions when cows are nursing calves, producers could consider early weaning instead of creep feeding calves to sustain a targeted weight gain. Creep feeding calves in drought conditions does not result in reduced nutrient needs of the dam. Data would suggest that creep-fed calves still nurse their dams so there is very little lactational and therefore nutrient pressure taken off the dam. In drought conditions, producers may want to consider early weaning the calves, especially if they have the facilities to feed lightweight calves, because they are very efficient at converting feed to weight gain. Information on this management alternative can be found in NebGuide G2047, *Management of Early Weaned Calves*. If cows are dry lotted in drought conditions, calves do not need to be creep fed as the calf will eat from the bunk alongside their dam. Make sure feed intake of the calf is accounted for in total pounds of feed delivered to the bunk. The calf will eat about 1% of its body weight on a dry matter basis.

Table I. Examples of ingredient compositions^a of creep feed.

	Creep 1	Creep 2	Creep 3	Creep 4	Creep 5	Creep 5
Ingredient	%, as-fed basis					
Cracked corn	45	50	25			50
Whole oats		50				
Dried distillers grains			25	25		50
Soybean meal	10				20	
Soybean hulls	45		50	70	75	
Molasses				5	5	
Nutrient Analysis						
Crude Protein, %	12.6	11.4	15.9	16.4	19.5	19.8
TDN, %	80	80	84	82	80	95

^aConsider adding an ionophore.

Nutrient Requirements and Ration Composition

The dam supplies part of the nutrients needed for calf growth through her milk. For spring-born calves, milk and vegetative forage should allow for adequate weight gain of the calf. As the calf gets older, the amount of milk produced by the dam decreases and it becomes more difficult to meet the calf's nutrient needs for additional weight gain. In addition, for spring calves, the quality of grass decreases in late summer and fall; again, there is usually insufficient protein or energy in the forage for optimum gain.

If it is decided to creep feed calves nursing their dam, the ration must be energy and protein dense. This is because the rumen of the calf is small, so the calf will not eat a large amount of feed. As an example, for a 220-pound calf to gain 2.0 pounds per day, it must consume about 7 pounds of feed (milk, forages, plus creep feed) on a dry

matter basis. The ration needs to supply 1.32 pounds of protein/day and 5.0 pounds of TDN/day (0.70 Mcal/lb NEM; 0.44 Mcal/lb NEg). Example creep rations are in *Table I*.

Not only must a creep ration be nutrient dense, but it must also be palatable. Most commercially available creep rations are pelleted and highly palatable. For creep rations that are not pelleted, the ration must be dust-free and uniformly mixed so the calf cannot sort out ration ingredients. If liquid feed ingredients are added to the creep feed, they must be included at a level that doesn't impede creep feed flow through the feeder. Rolling or coarsely cracking grains rather than fine grinding, helps reduce dust and will also reduce rumen upsets of home-mixed creep diets.

Simple mixtures of grain and protein supplement make satisfactory creep rations. Pelleted creep will have less feed waste. Bulky feeds such as oats or wheat bran should be included to protect against overeating. Both ground oats and bran are good feeds for getting calves started on creep feed.

Ionophores are feed additives that increase feed efficiency and average daily gain. These compounds also can be used to control coccidiosis. Examples of an energy creep ration and high protein creep ration are shown in *Table II*.

Feed Conversion of Different Types of Creep Feeds

Two types of creep feeds are used by producers—energy creep feeds and protein creep feeds. The most common is the high-energy creep.

Table II. Example Calculation to Determine the Value of Creep Feeding^a

	1 No Creep	2 Creep
A Weaning weight, lb.	500	565
B. Amount of creep feed fed per calf ^b , lb		520 (A2-A1) x 8
C. Sale calf price, \$/lb	2.10 ^c	2.06
D. Calf Value, \$	1,050	1,163.90 A x C
E. Value of added calf weight, \$/lb		1.75 (D2-D1)/(A2-A1)
F. Value of added gain, (\$)		113.75 (A2-A1) x E2
G. Creep feed price per ton delivered, \$		400
H. Creep feed price per pound delivered, \$		0.20 G2/2000 lb
I. Cost of creep feed fed, \$ per calf		104.00 B2 x H2
J. Return per head from creep feeding, \$		9.75 ^a F2 -I2

^aCalculations of costs and return include feed costs and does not include feeder, equipment, or labor costs.

^bUsed a 8:1 conversion of pounds of feed per pound of gain.

^cCreep fed calves weigh more but there is a "price slide" for beef calves and feeder calves where lighter weight calves have a price per pound advantage compared to heavier weight calves.

Data suggest that feed conversion varies for calves fed high-energy creep feeds. After removing the highs and lows for feed conversions for the high-energy creep, feed to gain is about 8:1 (range 15:1 to 6.8:1). This means 8 pounds of creep feed on a dry matter basis are required per pound of gain by the calf. Average daily gain for calves fed a high-energy creep feed is between 0.20 and 0.30 pound greater, compared to non-creep fed calves.

Calves fed the high-protein creep will eat less creep feed, have better feed conversions, and gain less total weight over the creep feeding period, compared to calves fed high-energy creeps. Data suggest that feed conversion for high-protein creep feeds on average is about 4:1, meaning that it takes 4 pounds of feed dry matter to produce a pound of gain.

Some of the wide variation in efficiency of feed use from creep feeding may be due to the following management factors:

- waste of feed due to wind, improper feeder design, or adjustment.
- location of the feeder, which affects frequency of eating and total feed intake.
- the creep feeder is not secure, and cows are able to consume some of the feed; and
- over-consumption of high-energy creep feeds.

Post-weaning Performance and Carcass Traits of Creep-Fed Calves

Generally, the greatest benefit of creep feeding is realized if calves are sold at weaning. There are conflicting data as to whether the added weight achieved using creep feeding is still realized at the end of the finishing period. There are data that suggest that non-creep fed calves will weigh similar to creep fed calves at the time of harvest. How calves are managed and fed after weaning would influence these weight differences. Calves that are managed for a low gain post weaning may not benefit from creep feeding, especially if they are fat. Again, creep feeding will benefit calves that go directly to the feedlot after weaning or sold within a few months of weaning. Creep fed calves will know how to eat out of a bunk and there will be less stress for these calves; therefore, less morbidity.

Finished calves that have been creep fed can have increased marbling at the time of harvest. Research suggests that marbling genes are influenced early in the calf's life, particularly in calves creep fed for at least 80 days. To retain

the response of creep feeding on marbling, calves need to be placed in the feedlot after weaning and fed a grain-based diet.

For Replacement Heifers

Heifers can be fed creep feed while nursing their dams. The challenge is when the additional gain is in the form of adding fat in addition to frame growth. Heifer calves can develop fatty tissue in the udder that consequently lowers milk production, compared to those heifers not fed high-energy creep rations. This impairment of future production can occur before the heifer reaches 3 months of age.

Creep feeding heifer calves while on their dams may be a management tool implemented in situations where the milk output of the dam is low and/or the growth potential is high. Creep feeding fall-born heifer calves may be warranted so they continue to grow at a rate to reach puberty by 13 to 15 months of age. Under most circumstances, it is usually easy to achieve the desired gain in heifer calves without the use of creep feeds. The possible exception could be for fall-born heifers.

Starting Calves on the Creep Ration

Calves often nibble at grain when they are about 3 weeks of age but will not eat appreciable amounts before they are 6 to 8 weeks old. If cows are grazing lush pasture and milking well, it is often difficult to get calves started on creep feed. Calves can be encouraged to start eating creep more quickly by:

- including palatable feed such as ground oats, wheat bran, cottonseed hulls, molasses.
- using an older calf that is already eating; or
- locating the feeder close to where cattle frequently congregate (water, shade, etc.).

Creep Feeder Design

Plans for constructing creep feeders for calves are published in MWPS-6, *Beef Housing and Equipment Handbook*, which is available at extension offices or online. The feeder should be roofed to protect feed from rain and should hold at least a week's supply of feed. This will depend on the size of the feeder and the number of calves it will accommodate. Allow 4 to 6 inches of trough space per calf. A feeder that is 8 feet long and feeds from both sides (16 feet of linear trough space) will accommodate 30 to 50 calves.

The feeder should be portable. If a fence is used around the feeder to keep the cows out, the entrance for the calves should be 1 foot 4 inches to 1 foot 6 inches wide by 3 feet high. Feeders not enclosed by a fence, or a perimeter barrier should be made strong enough to prevent cows and bulls from destroying the structure.

Location

For starting calves, locate the creep feeder near water or shade where cows rest frequently. Cows need to take calves to the creep until most calves have learned to eat. Once calves are eating, good pasture and range management dictates the creep be located away from water. Moving the feeder location frequently during the late summer and early fall may be desirable, especially in sandy soils to reduce vegetation damage. Large pasture or range situations may require more than one creep feeder for reasonable increases in rate and efficiency of gain from creep feeding.

For Starting Calves on Feed

Calves started on creep rations before weaning will begin eating sooner after they are weaned or shipped. They usually recover weaning or shipping weight losses in a shorter period and have lower morbidity and mortality rates, compared to calves that do not begin eating after being weaned. Some producers will creep feed calves while

they are still with their dam for a short period, two to four weeks before weaning. This practice is used to bunk-break the calves before weaning so at weaning they are eating.

Calculating Returns from Creep Feeding

Market price of heavier, fleshier feeder calves is usually lower than for calves that are lighter and have less condition. When cattle prices are high relative to feed prices, we can expect relative high price discrimination, as weights increase for calves of the same frame size and/or genetic potential. Only if cattle prices are low relative to feed prices may a premium exist for the heavier calves.

Net return calculations only consider the cost of the creep feed (*Table II*). Cost of feeders, equipment, and labor should be considered before initiating a creep feeding program.

These calculations should be completed before creep feeding to determine whether the management practice is cost-effective. For high-energy creep feed, use a feed to gain of 6:1 to 8:1, however, this varies and can be as high as 12:1.

The economics team at UNL created an easy-to-use Excel tool, “Creep Feeding cow-q-lator” or CFC. This tool can be accessed at the University of Nebraska-Lincoln webpage: <https://farmcents.unl.edu> and then clicking on the tab “Ag Manager Tools” where it is listed among other useful decision aids. This calculator will include equipment, labor and transportation costs in the calculations.

This publication has been peer reviewed.

**Index: Beef Cattle
Feeding and Nutrition**
Issued September 2024



This publication has been peer reviewed.
Nebraska Extension publications are available online
at <http://extension.unl.edu/publications>.

Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska-Lincoln cooperating with the Counties and the United States Department of Agriculture.

Nebraska Extension educational programs abide with the nondiscrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.

© 2024, The Board of Regents of the University of Nebraska on behalf of the University of Nebraska-Lincoln Extension. All rights reserved.